

**Table 4-2. Byssinosis: Mortality rates (per million population) by race and sex, U.S. residents age 15 and over, 1990-1999**

Year	Overall	White		Black		Other	
		Male	Female	Male	Female	Male	Female
<b>Crude Mortality Rate</b>							
1990	0.09	0.15	0.06	0.10	—	—	—
1991	0.06	0.09	0.05	0.10	—	—	—
1992	0.08	0.06	0.06	0.38	0.08	—	—
1993	0.08	0.09	0.06	0.37	0.08	—	—
1994	0.08	0.11	0.07	0.18	—	—	—
1995	0.07	0.13	0.03	—	—	—	—
1996	0.04	0.05	0.06	—	—	—	—
1997	0.05	0.09	0.01	0.09	—	—	—
1998	0.03	0.06	0.01	—	—	—	—
1999	0.03	0.03	0.04	—	—	—	—
<b>1990-1999</b>	<b>0.06</b>	<b>0.08</b>	<b>0.04</b>	<b>0.12</b>	<b>0.02</b>	—	—
<b>Age-Adjusted Mortality Rate</b>							
1990	0.09	0.17	0.05	0.14	—	—	—
1991	0.06	0.12	0.05	0.20	—	—	—
1992	0.07	0.06	0.05	0.46	0.11	—	—
1993	0.09	0.10	0.05	0.47	0.11	—	—
1994	0.09	0.13	0.06	0.39	—	—	—
1995	0.07	0.14	0.03	—	—	—	—
1996	0.04	0.05	0.05	—	—	—	—
1997	0.05	0.11	0.01	0.18	—	—	—
1998	0.03	0.07	0.01	—	—	—	—
1999	0.03	0.03	0.03	—	—	—	—
<b>1990-1999</b>	<b>0.06</b>	<b>0.10</b>	<b>0.04</b>	<b>0.18</b>	<b>0.02</b>	—	—

— indicates no deaths listed.

NOTE: See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause of death data. Population estimates from U.S. Bureau of the Census.